

Government of **Western Australia** Department of **Health** 



# Virus WAtch

## Week ending 23<sup>rd</sup> July 2023

### **Key Points**

#### Influenza and influenza-like illnesses (ILI)

- All indicators of influenza and ILI activity decreased in the past week including the rate of ILI presentations at sentinel GPs and at emergency departments (EDs).
- Influenza notifications to the Department of Health decreased but remained in the higher range of values usually reported at this time of year. Respiratory syncytial virus (RSV) notifications were comparable to the previous week.
- Non-influenza respiratory virus detections at PathWest Laboratory Medicine (PathWest) decreased in the past week.
- COVID-19 notifications decreased by 6% to 772 cases in the past week. See <u>COVID-19</u> <u>Weekly surveillance report (health.wa.gov.au)</u>

#### Gastroenteritis

- The rate of gastroenteritis presentations to sentinel GPs decreased in the past week while the rate of presentations to EDs increased.
- Rotavirus notifications to the Department of Health increased and norovirus detections at PathWest remained stable in the past week.


#### Other vaccine-preventable diseases

- **Chickenpox and shingles**: In the past week, the rate of chickenpox presentations to sentinel GPs increased and presentations to EDs remained stable. Shingles presentations to GPs increased and presentations to EDs decreased.
- Measles: No measles cases were notified in the past week.
- Mumps: No mumps cases was notified in the past week.
- Rubella: No rubella cases were notified in the past week.
- Invasive meningococcal disease (IMD): No IMD cases were notified in the past week.

#### Influenza and influenza-like illnesses (ILI)


The rate of ILI presentations to sentinel GPs continued to decrease in the past week (Figure 1).

Figure 1. Rate of ILI per 1000 consultations at sentinel GPs (Australian Sentinel Practices Research Network) in WA by week, 2018 to 2023 YTD




The rate of ILI-related presentations and admissions to EDs decreased in the past week (Figure 2).






Note: This graph is a count of current EDIS data using the ICD codes B34.9 and J06.9, which are consistent with a clinical presentation of influenza-like illness. This data may differ from that presented in the Winter Respiratory Illness Report provided by the Information and System Performance Directorate, DoH.

The number of respiratory illness presentations to EDs continued to decrease in the past week but remained in the higher range of values usually reported at this time of year (Figure 3).






Note: This graph is a count of current EDIS data using the ICD codes B34.9, H66.9, J00, J06.9, J09.0, J10.0, J10.1, J10.8, J11.0, J11.1, J11.8, J12.9, J18.0, J18.1, J18.8, J18.9, J20.9, J21.9, J22, J40, J44.0, J44.1, J44.9, J45.9, J46.0, J98.8, J98.9, R05 and COVID-19 code U07.1, which are consistent with a clinical presentation of all respiratory-like illness. This data is different to Figure 3 but similar to that presented in the Winter Respiratory Illness Report provided by the Information and System Performance Directorate, DoH.

In the past week, 640 influenza cases were notified to the Department of Health, which was 37% lower compared to the previous week (Figure 4).

Figure 4. Number of influenza notifications in WA by week, 2018 to 2023 YTD




Note: This graph is a count of all influenza notifications by week of receipt by the DoH, WA (through WANIDD) to the end of the current reporting week. The seasonal threshold defines a value above which may indicate seasonal influenza activity. The threshold value is calculated based on analysis of inter-seasonal influenza data from 2015 to 2018.

In the year to date, the number of influenza notifications and hospitalisations are higher than the previous five-year average, while the number of reported deaths is lower. Vaccination coverage to date is highest in the  $\geq$  65 year age group (Table 1). Thirty-three percent of influenza notifications have been in children under 10 years old (Figure 5).

# Table 1. Influenza notifications and vaccinationcoverage in WA, 2023 YTD

| Notifications   | Category                 | 2023 Year<br>to Date | 5 yr<br>average |
|---|--------------------------|----------------------|-----------------|
| Influenza infections<br>extracted by optimal<br>date of onset | Notifications            | 15,291               | 6,984           |
|   | Hospitalisations         | 1,710                | 1,072           |
|   | Reported<br>Deaths       | 5                    | 25              |
|   |                          | 2023 Year            | 5 yr            |
| Vaccinations  | Age group                | to Date              | average         |
|   |                          |                      |                 |
|   | 6mo-< 5 yrs              | 25.0%                | NA              |
| Influenza vaccinations  | 6mo-< 5 yrs<br>5-<15 yrs | 25.0%<br>18.0%       | NA<br>NA        |
| as recorded in the  |                          |                      |                 |
|   | 5-<15 yrs                | 18.0%                | NA              |


Figure 5. Influenza notifications by age group in WA, 2023 YTD



Note: Five-year average includes the years 2016 to 2019 and 2022. NA: data not available. Notification data source: WANIDD. Vaccination data source: AIR data downloaded from National Centre for Immunisation Research and Surveillance.

The seven-day moving average of influenza notification rates decreased in all regions except the Kimberley, where the rate remained stable (Figure 6).


Figure 6. 7-day moving average of influenza notifications per 100,000 people in WA by health region, 2023 YTD




Date of receipt

Note: This graph shows the 7-day moving average of influenza cases per 100,000 people in the WA health regions for 2023 by date of receipt, received by the DoH, WA (through WANIDD) to the end of the current reporting week.

The number of influenza cases reported as hospitalised decreased among adults and children in the past week (Figure 7).






Note: This graph shows the number of all notified influenza cases that have been hospitalised, by week of notification receipt, received by the DoH, WA (through WANIDD) to the end of the current reporting week. Child notifications were defined as individuals less than 18 years of age.


The influenza PCR test positivity decreased to 9% (134 detections) at PathWest in the last week. (Figure 8).






Note: This graph is a count of all WA samples reported by PathWest, excluding samples referred by other private laboratories for influenza subtyping.

PathWest reported 134 influenza detections in the past week; 70 (52%) were influenza A (which included 56 A/H1, 6 A/H3 and 8 influenza A not yet subtyped); 64 (48%) were influenza B (Figure 9). Of the total 640 influenza notifications in WA, 50% were influenza A and 48% were influenza B.






Note: The graph is a summary of all WA samples positive for influenza reported at PathWest, excluding samples referred by other private laboratories for influenza subtyping. These samples were tested using a rapid testing method that does not determine the influenza subtype (i.e., influenza A/H3N2 or A/H1N1).

In the past week, 560 respiratory syncytial virus (RSV) cases were notified in WA, which was comparable to the previous week (Figure 10).






Note: Respiratory syncytial virus (RSV) was made a notifiable infectious disease in WA in July 2021. This graph is a count of all RSV by week of onset by the DoH, WA (through WANIDD) to the end of the current reporting week.

Non-influenza respiratory virus detections at PathWest decreased in the past week (Figure 11). The most common non-influenza respiratory virus detected was RSV (169 cases).









Note: This graph is a count of all WA samples positive for a common respiratory virus other than influenza reported by PathWest.

In the past week, 772 COVID-19 cases were notified in WA, which was 6% lower compared to the previous week. Of these, 25% were diagnosed by PCR test and 75% were diagnosed by rapid antigen test (Figure 12).






Note: Data sourced from Public Health Operations COVID-19 Unified System (PHOCUS); Notification date is to the 6pm reporting period

The seven-day moving average of COVID-19 notifications per 100,000 population increased in the Great Southern, Midwest and Pilbara regions, remained stable in the Perth metropolitan area, and decreased in all other regions (Figure 13).






Note: Data sourced from Public Health Operations COVID-19 Unified System (PHOCUS). Western Australia Country Health Service (WACHS) region includes all nonmetropolitan health regions: Central-Wheatbelt, Goldfields, Great Southern, Kimberley, Midwest, Pilbara and South West. Perth metropolitan region includes East Metropolitan Health Service, North Metropolitan Health Service and South Metropolitan Health Service. Population denominator sourced from Australian Bureau of Statistics 2020 estimates. See also the <u>WA COVID-19 Weekly surveillance report</u> for further epidemiological analysis and the <u>Australian Government Dept of Health and Aged Care</u> for immunisation coverage data.

#### Gastroenteritis


The rate of gastroenteritis presentations to sentinel GPs decreased below the baseline in the past week (Figure 14).

Figure 14. Number of gastroenteritis presentations per 1000 consultations at sentinel GPs (Australian Sentinel Practices Research Network) in WA by week, 2018 to 2023 YTD




Gastroenteritis presentations at EDs increased in the past week and remained below the range of values usually reported at this time of year (Figure 15).





In the past week, statewide rotavirus notifications to the Department of Health increased and norovirus detections at PathWest remained stable (Figure 16).

Figure 16. Number of rotavirus notifications to the Department of Health and norovirus detections at PathWest in the past eight weeks




Note: Rotavirus notifications reported to the Department of Health include detections from all WA pathology laboratories. Norovirus detections are from PathWest only.

#### Viral rashes

The rate of chickenpox and shingles presentations to sentinel GPs increased above the baselines in the past week (Figure 17).


Figure 17. Number of varicella-zoster presentations per 1000 consultations at sentinel GPs (Australian Sentinel Practices Research Network) in WA in the past eight weeks



Note: Baseline levels for chickenpox and shingles presentations to WA ASPREN GPs per thousand consultations were calculated using the mean of weekly WA ASPREN data from week 1, 2018 to week 52, 2022.

Chickenpox presentations to EDs remained stable in the past week while shingles presentations decreased below the baseline (Figure 18).





Note: Baseline levels for varicella-zoster virus presentations to Emergency Departments in WA were calculated using the mean of weekly EDIS data from week 1, 2018 to week 52, 2022.

#### **Report Notes**

Virus WAtch is a weekly electronic publication by the Communicable Disease Control Directorate (CDCD) and key collaborators. It provides a brief summary of general practice and hospital emergency department sentinel surveillance data on influenza-like illness, gastroenteritis, and varicella-zoster disease, together with relevant laboratory information, to alert health care workers in WA about important circulating viruses. All figures and data were accurate at time of publication, but subject to change. Please note that the influenza and ILI surveillance systems in Western Australia (WA) have been impacted by the COVID-19 pandemic. Therefore, respiratory viral activity should be interpreted with caution and take into account the effects of changes in health seeking behaviour including accessing alternate health services such as telehealth, focused testing for COVID-19 at COVID-19 clinics or specific acute respiratory infection clinics, increased testing for other respiratory viruses and the impact of international border closures. The data collections used to create this publication include:

- Sentinel general practice (GP) data collected by WA members of the Australian Sentinel Practices Research Network (ASPREN).
- Emergency Department (ED) data provided by the Emergency Department Information System (EDIS), which currently incorporates data from the following hospitals: Fiona Stanley Hospital, Sir Charles Gardiner Hospital, Royal Perth Hospital, Perth Children's Hospital, King Edward Memorial Hospital, St John of God Midland, Bunbury Hospital, Armadale Hospital, Joondalup Health Campus, and Rockingham General Hospital.
- Disease notification data are sourced from the Western Australian Notifiable Infectious Diseases Database (WANIDD). These data are received by CDCD, WA Department of Health from medical providers and public or private laboratories in WA. Hospitalisation data are included in the report during the influenza season.
- Viral laboratory data obtained from PathWest laboratories at QEII Medical Centre, as well as via notification data sent by all WA laboratories to CDCD, WA Department of Health.
- As of 1 January 2022, the definition of a confirmed influenza case has changed to remove 'Single high titre by CFT or HAI to influenza virus' from the list of <u>laboratory definitive evidence</u>.
- As of March 2022, this report includes COVID-19 cases diagnosed by Polymerase Chain Reaction (PCR) test and Rapid Antigen Test (RAT) sourced from Public Health Operations COVID-19 Unified System (PHOCUS).
- Current and archived issues of Virus Watch <a href="http://ww2.health.wa.gov.au/Articles/F\_l/Infectious-disease-data/Virus-WAtch">http://ww2.health.wa.gov.au/Articles/F\_l/Infectious-disease-data/Virus-WAtch</a>.

#### This document can be made available in alternative formats on request for a person with disability.

© Department of Health 2023

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism, or review, as permitted under the provisions of the *Copyright Act 1968*, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.